

# Effect entomopathogenic fungi in control *Culex pipiens*

Mohammed Qasim Waheeb  
Dept. of Biology/College of Science/Al  
Muthanna University  
Samawah, 66001, Iraq  
E-mail: mhmdkas@mu.edu.iq

2<sup>nd</sup> Sadik Thajeb Ali  
Dept. of Biology/ College of  
Science/University of Thi-Qar  
Thi-Qar, 64001, Iraq

3<sup>rd</sup> Haider Radhi Malih  
Dept. of Biology/ College of  
Science/University of Thi-Qar  
Thi-Qar, 64001, Iraq

**Abstract**— *Culex pipiens* mosquitoes are thought to be vectors for many arboviruses, including West Nile virus and encephalitis virus, which have a global impact on human health. The natural management of this pest's aquatic stages is critical for sustaining an insecticide-free environment. The current study focused on the biological and biochemical effects of the entomopathogenic fungi *Beauveria bassiana* and *Trichoderma harizanium* on *Culex pipiens* laboratory colony 3rd instar larvae. The results showed that *B. bassiana* had the highest larval mortality (22%) with a concentration  $1 \times 10^6$  spore/ml and the period exposure recorded (36.6%) shortest lethal time (120 hrs), followed by *T. harizanium* (17.3%) with a concentration  $1 \times 10^7$  spore/ml and the period exposure recorded (29.1%) shortest lethal time (120 hrs), and had the lowest percent mortality (16%) with the concentration  $1 \times 10^3$  spore/ml and period exposure with (6%) longest (24 hrs) for *B. bassiana* and had the lowest percent mortality (4%) with the concentration  $1 \times 10^3$  spore/ml for *T. harizanium* and period exposure with 24 hrs (12.5%).

**Keywords**— *Culex pipiens*, *Beauveria bassiana*, *Trichoderma harizanium*, entomopathogenic fungi.

## I. INTRODUCTION

Mosquitoes are dipteran insects that serve as biological and mechanical vectors for a variety of parasites and pathogens that cause communicable diseases. They can transmit enzootic or even epizootic diseases like malaria, dengue fever, and filariasis, among others. The World Health Organization (WHO) created a Global Vector Control Response (GVCR, 2017-2030) to implement long-term vector control strategies (1). Despite its effectiveness, chemical control of vector insects poses a health, environmental, and climatic risk. The relevant situation of insecticide resistance and unsustainable interventions pose challenges to achieving sustainable development goals. The Culicidae are almost all bloodsuckers and are responsible for the spread of many serious diseases (2). The *Cx. pipiens* arouses the most interest due to its widespread distribution in tropical and subtropical countries, which has a socioeconomic impact. To counteract adult resurgence during adult control, vector control strategies typically target the aquatic stages of mosquitoes in their breeding habitat. Chemical larvicides that target mosquito breeding sites cause resistance in the targeted species as well as long-term secondary effects on non-targeted organisms, harming aquatic fauna (3). Exploring eco-friendly and biological control methods is necessary for developing an alternative

strategy for larval control. Entomopathogenic microorganisms hold a significant place among alternative methods of combating insect pests. The use of entomopathogenic fungi to control insect pests yielded promising results, with *Beauveria bassiana* (4) and *Metarhizium anisopliae* (5) being self-sustaining and efficient alternatives for controlling *Cx. pipiens* (6). Both species have a diverse range of isolates that differ in host specificity and origin. By controlling the aquatic stages of mosquitoes, these fungi will help to maintain the ecological balance in the surrounding aquatic habitats. (7). The current study examines the effectiveness of two Entomopathogenic microorganisms against *Cx. pipiens* larvae.

## II. METHODS

### A. Insect Colony Maintenance

A laboratory strain of *Cx. Pipiens* were obtained from the laboratory research insect of the Department of Biology, College of Science University of Thi-Qar, Iraq. The colony was kept in a walk-in chamber insectary at 27°C, 70% relative humidity, and a photoperiod of 12:12 hours. Mosquito larvae were grown in white enamel dishes with 1500 ml of distilled water. The newly hatched larvae were fed fish food (Tetra-Min, Germany). The adult was raised in wooden cages (24 x 24 x 24 cm) with 10% sucrose solution and a pigeon for female feeding.

### B. Fungus Culture

*Beauveria bassiana* (Balsamo), and *Trichoderma harizanium* isolates were obtained from the Mycology Center, Ministry of Science, and Technology, Iraq. The isolates were cultured in flasks on Sabouraud dextrose yeast agar (SDYA) medium (8), which contained 40 g glucose, 20 g peptone, 20 g agar, and 2 g yeast extract dissolved in 1000 ml of distilled water. The flasks were autoclaved for 15-20 minutes at 121°C. The media was poured into Petri dishes and ready for inoculation (9).

### C. Inoculum Preparations

Fungal cultures were plated into prepared Petri dishes and incubated in darkness at 25°C for 14 days. Conidial suspensions were made in a laminar airflow chamber by scraping cultures with a sterile inoculation needle and transferring them to 10 ml of distilled water containing 0.05% Tween 80. For 10 minutes, the mixture was stirred. The hyphal bodies were removed from the mixture by



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

filtering it through a fine mesh sieve. The final suspension's conidial concentration was determined by direct count with a hemocytometer (9, 10).

#### D. Bioassay

The purpose of the virulence test was to compare the efficacy of the two fungal isolates against *Cx. pipiens* 3rd larval instars. The fungal spore suspension was serially diluted in distilled water containing Tween-80 (0.1%) and stored at 5 °C until use (10). Conidia from the isolates were tested against larvae by adding the fungal suspensions to plastic cups containing 50 ml of distilled water and 25 3rd instar larvae. Each cup was inoculated with 1ml of fungal suspensions containing  $1 \times 10^6$ ,  $1 \times 10^7$ , and  $1 \times 10^8$  spores per milliliter. Control treatments included the addition of distilled water containing Tween-80 (0.1%) (11). Larvae were fed fish food and monitored daily. Mortality was measured at 24, 48, and 72 hours after larval feeding (12). The following biological parameters were studied to determine the effect of fungal treatment on the mosquito life cycle and female fecundity parameters: mean larval and pupal duration and percentage of pupation. Pupae were sexed and placed in glass globes in pairs. The percentages of adult emergence, adult longevity, female fecundity, and egg fertility were calculated (13).

#### E. Statistical analysis

One way ANOVA at  $p \leq 0.05$  was used to analyse the triplicate data means followed by a Multiple Comparison Test (MCT) to indicate the significant differences between means using a statistical analysis system (SAS, 2003).

### III. RESULTS

#### A. Effect suspension to the *Beauveria bassiana* and *Trichoderma harizanium* with death mortality on third larvae *Culex pipiens*

The figures (1-2) have been showing the effect of interface different concentrations from suspensions of *Beauveria bassiana* and *Trichoderma harizanium* fungi in death rate mortality on the fourth larvae stages for *Culex pipiens*, where exceeding the suspended *B. bassiana* fungus on the suspended *T. harizanium* fungus when used the most concentration  $1 \times 10^6$  spores/ml reached death mortality in the third stage of larvae *C. pipiens*, the highest concentration was registered mortality for both fungi at 22% and 17.3% respectively, while the period exposure for the same stage of larvae was mortality for the most period exposure as follow 36.6% and 29.1% respectively.




Figure (1) Effect suspension of the *B. bassiana* on larva stage (3) *Cx. pipiens*.

LSD 0.05(con.) =9.5, LSD 0.05 (p.ex.) =5.05, LSD 0.05 (interference)=7.76




Figure (2) Effect suspension of the *T. harizanium* on larva stage (3) *Cx. pipiens*.

LSD 0.05(con.) =7.83, LSD 0.05(p.ex.) =3.59, LSD 0.05 (interference.)=6.254.

### IV. DISCUSSION

The increasing in death rate with increase the concentration attributed to increased number of spores and then increased rate spores growth when attacked the host and impairment the immune system for on the confront for the insect, however the immune system for larvae can defend about body it, if the low concentrations, while the high concentrations may be loss the immune system effective it (14), and the statistical analysis has been proven presence significant differences between the percentage of the larvae stages and it has been proven between the species fungi, while the relationship between the death rate and the age of larvae stages reverse, the reason may be to the immune system is not completely, and as the body system is delicate and which allow penetration easily by the fungus spores (15), the reason different may be to the fungi disparity in death rate to ability these fungi secrete analytic enzymes and toxins effect on action lead to death it. (16), found the exposure of larvae of *Cx. quinquefasciatus* mosquitoes to *B. bassiana* fungus spores led to dead all it with concentration  $1 \times 10^8$  spores/ml and the death rate the percentage 97% with a concentration  $1 \times 10^7$  spores /ml showed a percentage of death rate *Cx. quinquefasciatus* mosquitoes increased with advance the suspension concentrations *Chrysosporium keratinophilum*, and not different about) the results that previously it was the percentage first stage of larvae for *Cx. quinquefasciatus* as follows (77.7, 60, 59.5) % with concentrations ( $2 \times 10^4$ ,  $2 \times 10^3$ ,  $2 \times 10^2$ .) spores/ml these result agreed with the results (17) when describing the relationship between the stages of larvae and for *Aedes aegypti* and death rate, where demonstrated death rate decrease with advancing the age and the death rate for the first and second stage reached 100% follow the third and second stage reached mortality 40% when the treatment with the suspension *Leptolegnia chapmani* with the concentration  $3.65 \times 10^5$  spores/ml. The (18) used treatment *An. stephensi* and *An. gambiae* with suspension *M. anisopliae* were the percentage mortality to the third and fourth stages less than the percentage mortality to the first and second stages for both species mosquitoes. The (19) showed to exposure the second stage of *Cx. quinquefasciatus* mosquito to the *A. niger* spores fungus with the concentration  $2 \times 10^6$  spores/ml led to death rate with 68.4% after two days from treatment, the injured insect by fungi may be lived the period progression

3-5 days because the grown spores and penetration the mycelium through the breathing tubes that caused suffocation the larvae because the closed breathing tubes, as well as the growth fungus in mid canal of trachea for larvae and drain the foods caused after 72 hours crash the lipid tissues and therefore may be percentage mortality to 100% after 96 hours, and also of some larvae dead during the metamorphosis (20). The (21) added the swallowing of larvae following secretion toxins from the entomofungal pathogen and thus leading toxemia(22).

#### V. CONCLUSIONS

Entomopathogenic fungi are thought to be a naturally occurring microbial control agent against many insects, helping to reduce host population in epizootics. The majority of them begin the infection process in the gut. Fungal enzymes aid in cuticle penetration, and toxin production induces the host's immune response, including activation/deactivation of host enzymes. The fungal propagation within the host body causes a significant depletion of host biomolecule availability, affecting variable parameters in the host life cycle, and eventually host death in a susceptible host. The difference in cuticular structure and chemical composition of the epicuticle influences the host-fungus relationship. Toxicity differences between selected fungal isolates.

#### REFERANCE

- 1- WHO, 2019. World Health Organization, "framework for the implementation of the global vector control response in the WHO African region", 2019.
- 2- X. Qin, X. Zhao, S. Huang, , J. Deng, X. Li, Z. Luo & Y. Zhang, "Pest management via endophytic colonization of tobacco seedlings by the insect fungal pathogen *Beauveria bassiana*". *Pest Management Science*, 77(4), 2007-2018,2021
- 3- S. O. Hamid, N. I. A, F. A. Halloran, T. M. A, F. Z. Bissaad & F. A. Benzina, "Study about the effect of *Beauveria bassiana* (Vuillemin IN 1912) on the aquatic stages of *Culex pipiens*" . (LINNÉ, 1758), 2013.
- 4- J. Ziani, "Application de *Beauveria bassiana* contre la punaise terne *Lygus lineolaris*"(Palisot de Beauvois)(Hémiptères: Miridés) dans les vignobles. Thesis2008.
- 5- O. Benserradj, I.Mihoubi, , "Larvicidal activity of entomopathogenic fungi *Metarhizium anisopliae* against mosquito larvae in Algeria". *Int. J. Curr. Microbiol. Appl. Sci* 3, 54-62,2014.
- 6- N.T. Loc, V.T.B.Chi, N.T.Nhan, , T.T.B.Hong N.T.P.Chi, N.T.Nghia, 2010. "Exploitation of *Beauveria bassiana* and *Metarhizium anisopliae* as potential biocontrol agents in integrated pest management (ipm) on citrus. Omonrice ",17, 152-163, 2010.
- 7- M. Farenhorst, B.G.Knols, . :Fungal entomopathogens for the control of adult mosquitoes". a look at the issues, Proceedings of the Netherlands Entomological Society meeting, pp. 51-59, 2007.
- 8- R. Sabouraud, , "Ann. Dermatol. Syphilol". 3:1061, 1982.
- 9- M.Osman, C.Stigloher, "Mueller, M.J., Waller, F., 2020. An improved growth medium for enhanced inoculum

production of the plant growth-promoting fungus *Serendipita indica*". *Plant methods* 16, 1-7,2020.

- 10- M. Alagawany, M. Madkour, M.T.El-Saadony, F.M. Reda, . "Paenibacillus polymyxa (LM31) as a new feed additive: Antioxidant and antimicrobial activity and its effects on growth, blood biochemistry, and intestinal bacterial populations of growing Japanese quail". *Anim. Feed Sci. Technol.* 276, 114920, 2021.
- 11- E.N.Haron, M.Ahmed, , S.S.Ali, A.A.Abas, M.E.Elshaier, "Evaluate the effects of entomopathogenic fungi isolates on Wheat Aphid, *Schizaphis graminu* (Rondani) (Hemiptera: Aphididae)". *Egypt. Acad. J. Biol. Sci. A, Entomology* 13, 149-159, 2020.
- 12- K.Javed, H.Javed, T. Mukhtar, D.Qiu, "Pathogenicity of some entomopathogenic fungal strains to green peach aphid, *Myzus persicae* Sulzer (Homoptera: Aphididae)." *Egypt J. Biol. Pest Control* 29, 1-7, 2019.
- 13- A.M. Saad, A.S.Mohamed, M.T. El-Saadony, M.Z.Sitohy, , "Palatable functional cucumber juices supplemented with polyphenols-rich herbal extracts". *LWT - Food Sci. Technol.* 148, 111668,2021.
- 14- E. J. Scholte, B. N. Njiru, , R. C. Smallegange W. Takken, & Knols, 2003 "Infection of malaria (*Anopheles gambiae* ss) and filariasis (*Culex quinquefasciatus*) vectors with the entomopathogenic fungus *Metarhizium anisopliae*". *Malaria journal*, 2(1), 1-8,2003.
- 15- F. E.Vega, & P. F.Dowd"The role of yeasts as insect endosymbionts. *Insect-fungal associations: ecology and evolution*", 211-243,2005.
- 16- G.Gayathri, C. Balasubramanian, P. V.Moorthi, & T. Kubendran, (2010). "Larvicidal potential of *Beauveria bassiana* (Balsamo) Vuillemin and *Paecilomyces fumosoroseus* (Wize,) Brown and Smith on *Culex quinquefasciatus* (Say)". *Journal of Biopesticides*, 3(1), 147,2010
- 17- T.Mc Innis Jr, & W. C. Zattau, "Experimental infection of mosquito larvae by a species of the aquatic fungus *Leptolegnia*. *Journal of Invertebrate Pathology*", 39(1), 98-104,1982.
- 18- T.Bukhari, A.Middelmann, C. J.Koenraadt, W.Takken, & B. G.Knols, "Factors affecting fungus-induced larval mortality in *Anopheles gambiae* and *Anopheles stephensi*". *Malaria journal*, 9(1), 1-15,2010.
- 19- K. Foo, J. S.Sathiya Seelan, & M. M .Dawood, "Microfungi associated with *Pteroptyx bearni* (Coleoptera: Lampyridae) eggs and larvae from Kawang River, Sabah (Northern Borneo)". *Insects*, 8(3), 66,2017.
- 20- S. T. Jaronski, ("Mass production of entomopathogenic fungi—state of the art". *Mass production of beneficial organisms*, 317-357,2023.
- 21- C. M.Lacey, L. A. Lacey, & , D. R. Roberts "Route of invasion and histopathology of *Metarhizium anisopliae* in *Culex quinquefasciatus*". *Journal of invertebrate pathology*, 52(1), 108-118, 1988.
- 22- S. M. Kazim,. "Effect of Muslin Bagging Technique on the Control of Greater Date Moth: *Arenipses*." *JOURNAL OF THI-QAR SCIENCE* 3, no. 2, 2012.